

ARE211, Fall 2009

CALCULUS2: THU, OCT 22, 2009

PRINTED: AUGUST 27, 2009

(LEC# 17)

CONTENTS

4. Univariate and Multivariate Differentiation (cont)	1
4.3. Partial, Directional and Total derivatives	2
4.3.1. Directional derivatives	2
4.3.2. Computing Directional Derivatives from Partial Derivatives	3
4.3.3. A nondifferentiable function whose partial derivatives exist	8
4.3.4. Total Derivative	9

4. UNIVARIATE AND MULTIVARIATE DIFFERENTIATION (CONT)

Key Points:

(1) Defn of a directional derivative: $f_h(\mathbf{x}) = \lim_{|k| \rightarrow \infty} \frac{(f(\mathbf{x}_0 + \mathbf{h}/k) - f(\mathbf{x}_0))}{\|\mathbf{h}\|/k}$ (equation (1) on p. 3).

(2) Defn of a differentiable function: $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *differentiable* at \mathbf{x}_0 if $\nabla f(\mathbf{x}_0)$ exists and if for all $\mathbf{h} \in \mathbb{R}^n$,

$$\lim_{|k| \rightarrow \infty} \frac{(f(\mathbf{x}_0 + \mathbf{h}/k) - f(\mathbf{x}_0)) - \nabla f(\mathbf{x}_0) \cdot \mathbf{h}/k}{\|\mathbf{h}\|/k} = 0 \quad (\text{equation (2) on p. 3})$$

or equivalently

$$\lim_{|k| \rightarrow \infty} \frac{(f(\mathbf{x}_0 + \mathbf{h}/k) - f(\mathbf{x}_0))}{\|\mathbf{h}\|/k} = \nabla f(\mathbf{x}_0) \cdot \frac{\mathbf{h}}{\|\mathbf{h}\|} \quad (\text{equation (3) on p. 5}).$$

- (3) A sufficient condition for differentiability of a function is that each of its partial derivatives is a continuous function.
- (4) Relationship between directional derivatives and total derivatives.

4.3. Partial, Directional and Total derivatives

4.3.1. *Directional derivatives.* Suppose we have a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and we want to know how it behaves as you move out in some direction starting from some given point in the domain \mathbf{x}^0 . In particular, want to know the slope of f in this direction.

Example: The directional derivative of f in the direction e_i —i.e., the vector whose i 'th component is 1, and others are zeros—is just the i 'th partial derivative of f .

Intuitively, imagine the graph of f is in fact a cake. Now “cut” the cake in the direction we’re interested in, so that the cut passes through \mathbf{x}^0 . Look at the cross-section of the cake you’ve obtained. It’s just like the graph of a function from \mathbb{R} to \mathbb{R} , and, if the graph is smooth, it has a slope. That slope is going to be the directional derivative in the direction you’ve selected. Having said that, it’s not so straightforward to compute it. For computational purposes, it’s useful to extrapolate from the following (obvious) facts:

- if $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable, then the slope of f at x equal to the differential of f at x , evaluated at 1, i.e., $f'(x) = df^x(1)$.
- if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable, then the i 'th partial derivative of f at \mathbf{x} is equal to the differential of f at \mathbf{x} , evaluated at \mathbf{e}^i , the i 'th unit vector, i.e., $e_k^i = \begin{cases} 1 & \text{if } k = i \\ 0 & \text{if } k \neq i \end{cases}$, i.e., $f_i(x) = df^x(\mathbf{e}^i)$.
- by analogy if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable, then the directional derivative of f at \mathbf{x} in an arbitrary direction h had better be equal to the differential of f at \mathbf{x} , evaluated at the (unique) *unit length* vector that points in the direction h . If this weren't true, then partial derivatives would not be special cases of directional derivatives!

This observation motivates why there's an $\|\mathbf{h}\|$ in the denominator of the definition below: note that if the h in the numerator is of unit length, then that term disappears, if not, the norm makes the necessary adjustment.

Definition: Given $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $\mathbf{h} \in \mathbb{R}^n$, the directional derivative of f at \mathbf{x}_0 in the direction \mathbf{h} is given by¹

$$\lim_{|k| \rightarrow \infty} \frac{(f(\mathbf{x}_0 + \mathbf{h}/k) - f(\mathbf{x}_0))}{\|\mathbf{h}\|/k} \quad (1)$$

Notice that the directional derivative of f at \mathbf{x}_0 in the direction \mathbf{h} has the same magnitude but the opposite sign from the directional derivative of f at \mathbf{x}_0 in the direction $-\mathbf{h}$.

The notation in (1) is shorthand for the following idea: there exists some scalar $c \in \mathbb{R}$ such that for *any* sequence, $\{k(n)\}$ with the property that $|k(n)|$ increases without bound with n , the expression in (1) must converge to c . To determine whether a directional derivative exists, it is instructive *but not sufficient* to consider whether the limit (1) exists for the particular sequence $k(n) = n$. You also have to worry about sequences that approach \mathbf{x}_0 from the opposite direction, e.g., $k(n) = -n$. Fortunately, you virtually never have to compute the value of limit c (or if such a limit exists) by considering *specific* sequences. Typically you do as the example does on p. 8, which is to make some terms disappear because some power of k is in the denominator.

4.3.2. Computing Directional Derivatives from Partial Derivatives. Computing directional derivatives can be a huge pain. It turns out, however, that provided the function f is *differentiable*, you can infer any directional derivative just by knowing the partial derivatives of f . Specifically:

Definition: a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be *differentiable* at \mathbf{x}_0 if $\nabla f(\mathbf{x}_0)$ exists and if for all $\mathbf{h} \in \mathbb{R}^n$,

$$\lim_{|k| \rightarrow \infty} \frac{(f(\mathbf{x}_0 + \mathbf{h}/k) - f(\mathbf{x}_0)) - \nabla f(\mathbf{x}_0) \cdot \mathbf{h}/k}{\|\mathbf{h}\|/k} = 0 \quad (2)$$

¹ An alternative defn is $\lim_{|k| \rightarrow \infty} \frac{(f(\mathbf{x}_0 + \mathbf{h}/(k\|\mathbf{h}\|)) - f(\mathbf{x}_0))}{1/k}$. The difference is that in this alternative defn, the norm of \mathbf{h} is moved into the numerator. For a proof that the two definitions are equivalent, see </home/simon/teaching/classes/mathLectures/AltDirectionalDerivDefn.tex>




FIGURE 1. Flat board against the graph

Note that $(\mathbf{h}/k)_{k=1}^{\infty}$ is a sequence of vectors, all pointing in the same direction, whose lengths shrink to zero. Also, the sequence of k 's can change sign, just as in the primitive definition of the derivative of a function defined on \mathbb{R} .

This definition says, literally, that a function is differentiable at \mathbf{x}_0 if you can put a flat board up against the graph of the function, so that it touches the graph vertically above/below \mathbf{x}_0 , *and if nearby*, the board is very close to the graph (cf what happens when you put a board on top of a Hershey Kiss). Fig. 1 illustrates how you put a flat board against a graph. Note however, that the relationship between the graph and the board may be less pretty than in this figure: the board could be above the graph on one side of \mathbf{x}_0 and below it on the other side.² The important thing is that the board and the graph are *very close* in a neighborhood of \mathbf{x}_0 . To get a feel for what it means for the board to be close to the graph, try putting a board near the graph in Fig. 3 below, vertically above zero. You just can't do it.

² The board is a first order approximation to the function. Whether the board is above the graph, below it, or cuts through it depends on the definiteness of the Hessian. If the Hessian is positive (negative) definite, the board will be above (below) the graph. If it is indefinite, then the board will actually cut through the graph. If it is semi-definite, you have to check the third-order terms in the Taylor expansion.

Since the second term on the left hand side of (2), i.e., $\frac{\nabla f(\mathbf{x}_0) \cdot \mathbf{h}/k}{\|\mathbf{h}\|/k}$, clearly does not depend on k , we can rewrite definition (2) as:

$$\lim_{|k| \rightarrow \infty} \frac{(f(\mathbf{x}_0 + \mathbf{h}/k) - f(\mathbf{x}_0))}{\|\mathbf{h}\|/k} = \nabla f(\mathbf{x}_0) \cdot \frac{\mathbf{h}}{\|\mathbf{h}\|} \quad (3)$$

Since the length of the vector $\frac{\mathbf{h}}{\|\mathbf{h}\|}$ is unity, a verbal version of this definition is: f is differentiable at x_0 if for every direction \mathbf{h} , the directional derivative of f at x_0 in the direction \mathbf{h} , i.e., $\lim_{k \rightarrow \infty} \frac{(f(\mathbf{x}_0 + \mathbf{h}/k) - f(\mathbf{x}_0))}{\|\mathbf{h}\|}$ is equal to the value of the differential function, when evaluated at *the unique unit length* vector pointing in the direction \mathbf{h} . Some times in class I have sloppily talked as if the vectors \mathbf{h} and $-\mathbf{h}$ “point in the same direction,” i.e., that all that matters is the angle of the line thru the origin on which \mathbf{h} lies, and this line stays the same when you flip \mathbf{h} through 180 degrees. This was bad sloppy talk, as this definition clearly indicates. From now on, I'll try to consistently say that \mathbf{h} and $-\mathbf{h}$ “point in opposite directions.”

There is yet another way of saying what differentiability means, this time in terms of linear algebra. Loosely, a function is differentiable at \mathbf{x}_0 if the directional derivatives of f at \mathbf{x}_0 live in the *vector space* spanned by the partial derivatives. This statement is not precise: the following theorem makes it correct. Recalling the definition above of \mathbf{e}^i , the unit vector pointing in the i 'th direction, we have

Theorem: a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable at \mathbf{x}_0 iff $\nabla f(\mathbf{x}_0)$ exists and for every $\mathbf{h} \in \mathbb{R}^n$ with $\|\mathbf{h}\| = 1$, the pair $(\mathbf{h}, f_{\mathbf{h}}(\mathbf{x}_0))$ belongs to the span of the set of vectors $\{(\mathbf{e}^i, f_i(\mathbf{x}_0)) : i = 1, \dots, n\}$.

(Note that $(\mathbf{h}, f_{\mathbf{h}}(\mathbf{x}_0))$ and $(\mathbf{e}^i, f_i(\mathbf{x}_0))$ both live in $\mathbb{R}^n \times \mathbb{R} = \mathbb{R}^{n+1}$.)

Letting $I = [\mathbf{e}^1, \dots, \mathbf{e}^i, \dots, \mathbf{e}^n]$ denote the $n \times n$ identity matrix, the above theorem is saying that f is differentiable at \mathbf{x}_0 iff for every $\mathbf{h} \in \mathbb{R}^n$ with $\|\mathbf{h}\| = 1$,

$$\begin{bmatrix} h_1 \\ h_2 \\ \vdots \\ h_n \\ f_i(\mathbf{x}_0) \end{bmatrix} = h_1 \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ f_1(\mathbf{x}_0) \end{bmatrix} + h_2 \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ f_2(\mathbf{x}_0) \end{bmatrix} + \cdots + h_n \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \\ f_n(\mathbf{x}_0) \end{bmatrix}$$

Thus, this theorem simply restates the definition of differentiability in vector space language. The span of the set of vectors $\{(\mathbf{e}^i, f_i(\mathbf{x}_0)) : i = 1, \dots, n\}$ is precisely the tangent plane, translated back to the origin. The theorem says that a function is differentiable iff for each direction \mathbf{h} in the unit circle, the directional derivative $f_{\mathbf{h}}(\mathbf{x}_0)$ is the height of the tangent plane vertically above \mathbf{h} .

Take an example: $f(\mathbf{x}) = x_1x_2$, $\bar{\mathbf{x}} = (1, 2)$, $\nabla f(\mathbf{x}) = (x_2, x_1)$, $\nabla f(\bar{\mathbf{x}}) = (2, 1)$, $\mathbf{h} = (\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$. Now

$$\begin{aligned} \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ f_i(\bar{\mathbf{x}}) \end{bmatrix} &= h_1 \begin{bmatrix} 1 \\ 0 \\ f_1(\bar{\mathbf{x}}) \end{bmatrix} + h_2 \begin{bmatrix} 0 \\ 1 \\ f_2(\mathbf{x}_0) \end{bmatrix} \\ &= \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} + \frac{1}{\sqrt{2}} \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \\ \frac{3}{\sqrt{2}} \end{bmatrix} \end{aligned}$$

On the other hand, consider the function graphed in Fig. 3 below. At the origin, both partials for this function are zero, so that span of $\{(\mathbf{e}^i, f_i(\mathbf{0})) : i = 1, 2\}$ is the horizontal plane. However, almost none of the $(\mathbf{h}, f_{\mathbf{h}}(\mathbf{0}))$'s live in this plane.

So far, we have been talking about differentiability *at a point*. We now define what it means for a *function* to be differentiable.

Definition: a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be *differentiable* if it is differentiable at every point in its domain.

Simon & Blume make a particularly egregious mistake when they say what differentiability is. They don't have a formal definition, but their very sloppy choice of words imply that a function is *differentiable* at \mathbf{x}_0 if $\nabla f(\mathbf{x}_0)$ exists. They rarely make egregious mistakes, but in this case... This illustrates the perils of using friendly wordy definitions (something I, of course, would *never* do) rather than writing out the math.

Obviously, it could be a big pain to check that for every unit length vector \mathbf{h} , the differential of f at \mathbf{x}_0 , evaluated at \mathbf{h} , coincides with the directional derivative of f at \mathbf{x}_0 in the direction \mathbf{h} . Fortunately, we don't have to do this, because of the following sufficiency theorem.

Theorem: Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$. Suppose that all partial derivatives of f exist and are continuous in a neighborhood of $\mathbf{x}_0 \in \mathbb{R}^n$. Then f is differentiable at \mathbf{x}_0 .

A function f defined on an open set U is said to be *continuously differentiable* if $\nabla f(\cdot)$ is a continuous function on U . (Notice that there's an abuse of terminology here: the whole point of the definition of differentiability is that *it's about much more than just the partials*. Yet continuous differentiability, a more stringent concept, concerns *only* the partials!) The above theorem establishes that continuous differentiability is a sufficient condition for differentiability. It is not, however, a necessary condition

for differentiability. To see this, consider the function $f(\mathbf{x}) = \begin{cases} x^2 \sin(1/x) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$. As Fig. 2

suggests, the graph of $f(\cdot)$ (plotted the left panel) has a tangent plane at zero—you have to use your imagination a bit to visualize it—but the derivative $f(\cdot)$ (given by $f(x) = 2x \sin(1/x) - \cos(1/x)$ and plotted in the right panel) oscillates increasingly wildly as you approach zero, and so is not a continuous function at zero. The set of continuously differentiable functions is denoted by \mathbb{C}^1 . More generally, the set of i -times continuously differentiable functions (i.e., each of the first partials are continuously differentiable, etc., etc) is denoted by \mathbb{C}^i .

Returning to the relationship between directional derivatives and the gradient for well-behaved functions, the following example illustrates that provided a function is differentiable, then computing a directional derivative in the primitive way—i.e., the left hand side of (3)—will yield the same answer as computing it as a linear combination of partials, i.e., the right hand side of (3).




FIGURE 2. A differentiable function that is not continuously differentiable

Example: Let $f = x_1x_2$, $\mathbf{x}_0 = \mathbf{h} = (1, 1)$. We'll compute the directional derivative of f at \mathbf{x}_0 in the direction \mathbf{h} , both ways:

(1) The primitive way: (left hand side of (3))

$$\lim_{|k| \rightarrow \infty} \frac{f(\mathbf{x}_0 + \mathbf{h}/k) - f(\mathbf{x}_0)}{\|\mathbf{h}\|/k} = \lim_{|k| \rightarrow \infty} \frac{\left(1 + \frac{1}{k}\right)^2 - 1^2}{\|(1, 1)\|/k} = \lim_{|k| \rightarrow \infty} \frac{\left(1 + \frac{2}{k} + \frac{1}{k^2} - 1\right)}{\sqrt{2}/k} = \frac{2}{\sqrt{2}}$$

(2) The indirect way: (right hand side of (3))

$$\nabla f(\mathbf{x}_0) \cdot \frac{\mathbf{h}}{\|\mathbf{h}\|} = \frac{\begin{bmatrix} 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 \end{bmatrix}'}{\sqrt{2}} = \frac{2}{\sqrt{2}}$$

4.3.3. *A nondifferentiable function whose partial derivatives exist.* Consider the function

$$f(x, y) = -\gamma(x^2 + y^2) + \begin{cases} \frac{xy}{\operatorname{sgn}(x-y)\sqrt{|x^2-y^2|}} & \text{if } |x| \neq |y| \\ 0 & \text{otherwise} \end{cases}$$

where $\gamma > 0$, is graphed in the top panel of Fig. 3. ($\operatorname{sgn}(\cdot)$ is a function that is 1 if the argument is positive, -1 if negative and zero if zero.) Look at the function in a neighborhood of zero. The function is continuous at zero, indeed, has beautifully behaved cross sections in *every* direction,

which are strictly concave in every direction (a typical cross-section is graphed in the bottom panel of Fig. 3). In particular, the partials w.r.t. x and y both exist and $\nabla f(\mathbf{0}) = \mathbf{0}$. To see this, note that for all $\alpha \neq 0$, $f(\alpha, 0) = f(0, \alpha) = 0/\alpha = 0$. Hence $\lim_{|x| \rightarrow 0} \frac{(f(\mathbf{0}+(x,0))-f(\mathbf{0}))}{\|\mathbf{x}\|} = \lim_{|y| \rightarrow 0} \frac{(f(\mathbf{0}+(0,y))-f(\mathbf{0}))}{\|\mathbf{y}\|} = 0$. However, as Fig. 3 indicates, these partials provide no information about what the slope of f is when you move in any direction other than parallel to an axis. To see that the requirement for differentiability fails, note first from the graph that you clearly can't put a tangent plane on top of the graph at $(0,0)$. More precisely, note that the directional derivatives *don't* live in the vector space spanned by the the vectors $\{(\mathbf{e}^i, f_i(\mathbf{x}_0)) : i = 1, \dots, n\}$. Also, observe that the *sufficient* condition for differentiability fails, i.e., neither partial is a continuous function in a neighborhood of zero. (The first partial is plotted in Fig. 4, where the domain is a circle of small radius centered at zero.)

4.3.4. *Total Derivative.* This concept appears to have been invented by economists but never appears in a math book. With good reason. It's not really a derivative at all; rather it's nothing more than a straightforward application of the chain rule. The *only* reason why it is taking up space in these notes is because economists have been using it for years, so you need to be familiar with it. It would be better to erase it from your brain (after this lecture), or if you want to leave it in your brain, then at least give it another name, which does not involve the word 'derivative'.

In any event, the total derivative should not to be confused with the total differential. Often, the arguments of a function depend on each other: we have a function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ which depends on x and w , but w itself depends on x , so we write $f(x, w(x))$. If we want to know how f will change when x changes, we need to take into account that w will change too.

General defn: The *total* derivative of $f : \mathbb{R}^n \rightarrow \mathbb{R}$ with respect to x_i is written $\frac{df(x)}{dx_i}$ (cf the partial derivative sign " ∂ ") and is defined by

$$\frac{df(\mathbf{x})}{dx_i} = f_1(\mathbf{x}) \frac{\partial x_1}{\partial x_i} + \dots + f_i(\mathbf{x}) + \dots + f_n(\mathbf{x}) \frac{\partial x_n}{\partial x_i}$$

Economic example: the firm's profit function $\pi(p, q(p)) = pq(p)$, where $q(p)$ is the optimal output choice given p . In this case, $\frac{d\pi(p, q(p))}{dp} = \frac{\partial \pi(p, q(p))}{\partial p} + \frac{\partial \pi(p, q(p))}{\partial q} \frac{dq}{dp} = q(p) + pq'(p)$.




FIGURE 3. Graph of f : partials convey no information about other directional derivatives

Total derivative, directional derivative and differential: What's the relationship between these three concepts? In the good old days, pre-enlightenment, I used to tell people that the total derivative




FIGURE 4. Graph of $f_1(\cdot)$: domain is a small circle around zero

was the *directional derivative* in the direction $(dp, q'(p)dp)$. I was **wrong**. The following reasoning explains why what I said was wrong:

- the *total derivative* of $\pi(\bar{p}, q(\bar{p}))$ tells you how much π changes when you increase p by *one unit*.
- the *directional derivative* of $\pi(\bar{p}, q(\bar{p}))$ in the direction $(dp, q'(p)dp)$ tells you how much π changes when you move *one unit in length* from $(\bar{p}, q(\bar{p}))$ in the direction $(dp, q'(p)dp)$.
- but if you increase p by one unit, you *don't* move *one unit* of length in the direction $(dp, q'(p)dp)$; in fact, you move $\|(1, q'(p))\|$ units of length in this direction!
- hence the *total derivative* of $\pi(\bar{p}, q(\bar{p}))$ is the differential of π at $(\bar{p}, q(\bar{p}))$, evaluated at the magnitude of the change, i.e., at $(1, q'(p))$.
- alternatively, the total derivative is the directional derivative in the direction $(1, q'(p))$, *multiplied by the length of the change*, i.e., by $\|(1, q'(p))\|$.

Example: The following example illustrates the above relationships. Consider a competitive industry with supply function $q(p) = \sqrt{p}$ and zero costs. (Economics is a little wierd, but this is a math course after all...) Industry profits are $\pi(p, q) = pq(p)$. Set $p = 1$ and note that $q(p) = 1$ while $q'(p) = 0.5/\sqrt{p} = 0.5$. Note also that $\nabla\pi(1, 1) = (q, p) = (1, 1)$. All four of the routes below tell you how much π goes up when you increase p by one unit, and q then increases by $q'(p)$. Happily all four routes give you the same answer.

(1) The derivative of π , viewed as a function of p only:

$$\left. \frac{d\pi(p, q(p))}{dp} \right|_{p=1} = \left. \frac{d}{dp} (p\sqrt{p}) \right|_{p=1} = \left(\sqrt{p} + \frac{0.5p}{\sqrt{p}} \right) \Big|_{p=1} = 1.5$$

(2) The total derivative of π , viewed as a function of p and $q(p)$:

$$\frac{d\pi(p, q(p))}{dp} = \frac{\partial\pi(p, q(p))}{\partial p} + \frac{\partial\pi(p, q(p))}{\partial q} \frac{dq}{dp} = 1 + 1 \star 0.5 = 1.5;$$

(3) The differential of $\pi(p, q)$ at $(1, q(1))$, evaluated at $\mathbf{h} = (1, q'(1))$:

$$\nabla\pi(1, 1) \cdot \mathbf{h} = \begin{bmatrix} 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0.5 \end{bmatrix}' = 1.5;$$

(4) The directional derivative of $\pi(p, q)$ in the direction $\mathbf{h} = (1, q'(p))$:

$$\nabla\pi(1, 1) \cdot \frac{\mathbf{h}}{\|\mathbf{h}\|} = (q(p), p) \cdot \frac{\mathbf{h}}{\|\mathbf{h}\|} = \frac{\begin{bmatrix} 1 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0.5 \end{bmatrix}'}{\sqrt{1.25}} = \frac{1.5}{\sqrt{1.25}}$$

Now if you add to $(1, 1)$ a vector pointing in the direction \mathbf{h} of length $\sqrt{1.25}$, you increase π by

$$\frac{1.5}{\sqrt{1.25}} \sqrt{1.25} = 1.5;$$